

The Angiographic Diagnosis of Gastrointestinal Bleeding

Diagnosis of the site of gastrointestinal hemorrhage is often difficult. The usefulness of selective arteriography in locating the source of gastrointestinal bleeding has been demonstrated. Bleeding as slow as 0.5 ml per minute has been localized in experimental animals. Selective angiography is particularly useful in determining bleeding sites in sections of the gastrointestinal tract which are difficult to endoscope—the small intestines and colon. In 27 patients with acute rectal hemorrhage and noncontributory sigmoidoscopy, selective angiography demonstrated a bleeding point in 18. In 13 patients, the hemorrhage was secondary to diverticulosis and in 12 of the 13 the bleeding was located to the right of the splenic flexure. Precise localization of the offending diverticulum allowed the surgeon to perform segmental colon resection. Selective arteriography is often the only effective way of diagnosing the source of chronic colonic and small intestinal bleeding caused by arterial-ve-

nous malformations. In one study three-quarters of the patients had a laparotomy which failed to reveal A-v malformations that were subsequently viewed when selective angiography was used. Bleeding from the second and third parts of the duodenum is difficult to localize by endoscopy and barium roentgenography, but bleeding stress ulcers in the mid-distal duodenum may be readily localized angiographically.

The value of selective catheterization in the bleeding patient with varices is two-fold. Angiography may demonstrate a nonvaricocoe source of hemorrhage, such as an ulcer. If no source of arterial hemorrhage is found, and portal hypertension with collateral varicocoe channels is demonstrated, selective vasopressin infusion through the arterial catheter will often control the active varicocoe hemorrhage.

GEORGE S. HARELL, M.D.

RONALD A. CASTELLINO, M.D.

REFERENCES

- Casarella WJ, Kanter IE, Seaman WB: Right-sided colonic diverticula as a cause of acute rectal hemorrhage. *N Engl J Med* 286:450-453, Mar 1972
- Alfidi RJ, Caldwell DE, Tarar R, et al: Recognition and angio-surgical detection of arteriovenous malformations of the bowel. *Ann Surg* 174:573-581, Oct 1971
- Baum S, Ward S, Nusbaum M: Stress bleeding from the mid-duodenum—An often unrecognized source of gastrointestinal hemorrhage. *Radiology* 95:595-601, Jun 1970

ADVISORY PANEL TO THE SECTION ON RADIOLOGY

GABRIEL WILSON, M.D., *Chairman*
Los Angeles
University of California, Los Angeles

R. THOMAS BERGERON, M.D.
Los Angeles County/
University of Southern California Medical Center
CMA Section on Radiology
Assistant Secretary

DUANE E. BLICKENSTAFF, M.D.
CMA Section on Radiology,
Secretary
La Jolla

RONALD A. CASTELLINO, M.D.
Stanford University

GLADDEN V. ELLIOT, M.D.
California Society of Radiologists,
President
La Mesa

VERNON R. GEE, M.D.
California Society of Radiologists
Redding

W. N. HANAFEE, M.D.
University of California, Los Angeles

GEORGE JACOBSON, M.D.
Los Angeles County/
University of Southern California Medical Center

MELVIN P. JUDKINS, M.D.
Loma Linda University

ELLIOT LASSER, M.D.
University of California,
San Diego

A. R. MARGULIS, M.D.
University of California,
San Francisco

BERNARD J. O'LOUGHLIN, M.D.
University of California,
Irvine,
California College of Medicine

ANTOLIN RAVENTOS, M.D.
University of California,
Davis

WARREN M. RUSSELL, M.D.
San Francisco

E. NICHOLAS SARGENT, M.D.
Los Angeles County/
University of Southern California Medical Center

JUSTIN J. STEIN, M.D.
CMA Scientific Board
Los Angeles